

INTERNSHIP PROPOSAL

Laboratory name: Institut des NanoSciences de Paris

CNRS identification code: UMR 7588

Internship director'surname: Agnès Maître / Fengzhi Gu

e-mail: agnes.maitre@insp.jussieu.fr

Phone number: 0144274217

Web page: [quantum nanophotonics](http://quantum.nanophotonics)

Internship location: Sorbonne université, Jussieu, 22-32, 5th floor

Thesis possibility after internship: YES

Funding: YES If YES, which type of funding: doctoral school

Strong coupling for a colloidal quantum dot inside a plasmonic nano-antenna

Keywords: *fluorescence, quantum dot, plasmonics, strong coupling*

CdSe/CdS core/shell quantum dots are excellent bright nano-emitters. At room temperature, under low excitation power they behave as high quantum efficiency single photon source thanks to the electronic confinement and efficient Auger processes leading to the recombination of a single exciton. They can be analysed as a simple two-level system and their emission linewidth is approximately $\Delta\lambda \approx 20$ nm. However, by increasing the excitation power, emission spectrum broadens dramatically up to $\Delta\lambda \approx 150$ nm (Fig1a). Moreover, the emission intensity grows non-linearly with increasing excitation power [1][2]. The two-level system paradigm fails for interpreting those features. In this view, we have successfully developed a model based on the radiative recombination of multiple excitonic levels within a single nanocrystal, relying on statistical description[3] of electron and hole populations in a quasi-equilibrium and Bayesian analysis.

Patch plasmonic nanoantennas consist of a thin dielectric medium sandwiched between a thick gold layer and gold patch. In the preceding years we have developed protocols in order to locate single quantum dots inside antenna. We have demonstrated that for a quantum dot inside such antenna, the absorption cross section is increased by 3 order of magnitude. Therefore such an antenna/quantum dot system is a good candidate for achieving strong coupling, a regime for which quantum dots emission and cavity resonance cannot be considered independently anymore.

During the internship, the student will start developing bright optical nanosources by coupling nanoemitters to plasmonic antennas. He/she will fabricate then and study by confocal optical microscopy the fluorescence (emission rate and spectrum, directivity,...). The final objective is to achieve the regime of strong coupling regime between the emitter and the confined field excited inside the antenna. Thanks to a high confinement and excitation, the emitters should gain specific original quantum properties.




Figure: 1) TEM images of CdSe/CdS nanocrystals 2) patch antenna 3) SEM image of patch antenna 4) emission pattern of patch antenna

References

[1] A.R Dhawan, et al. (2020) Light: Science & Applications, 9(1), 1-9., [2] A. R Dhawan et al (2022), Advanced Materials, 34(11), 2108120, [3] P. Wurfel. (1982) 15(18), 3967.

M1 and M2

Condensed Matter Physics: YES Quantum Physics: YES

Quantum Engineering: YES